

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/92199>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

pmoA Primers for Detection of Anaerobic Methanotrophs[▽]

Francisca A. Luesken,¹§ Baoli Zhu,^{1,2}§ Theo A. van Alen,¹§ Margaret K. Butler,¹†
Marina Rodriguez Diaz,¹‡ Bongkeun Song,³ Huub J. M. Op den Camp,¹
Mike S. M. Jetten,¹ and Katharina F. Ettwig^{1*}

Department of Microbiology, Institute for Water and Wetland Research, Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, Netherlands¹; State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China²; and Center for Marine Science, University of North Carolina Wilmington, Wilmington, North Carolina 28409³

Received 18 December 2010/Accepted 23 March 2011

Published *pmoA* primers do not match the *pmoA* sequence of “*Candidatus Methyloirabilis oxyfera*,” a bacterium that performs nitrite-dependent anaerobic methane oxidation. Therefore, new *pmoA* primers for the detection of “*Ca. Methyloirabilis oxyfera*”-like methanotrophs were developed and successfully tested on freshwater samples from different habitats. These primers expand existing molecular tools for the study of methanotrophs in the environment.

Aerobic methane oxidation is performed by several specialized groups of bacteria that are widely distributed in the environment (6). The first step of aerobic methane oxidation is the conversion of methane to methanol, catalyzed by either soluble or particulate methane monooxygenase (pMMO). The alpha subunit of pMMO, encoded by the *pmoA* gene, is highly conserved (5) and often used as a functional marker for analyzing methanotrophs in the environment (14). The first oligonucleotide primers targeting the *pmoA* gene, A189 and A682, have been used extensively in environmental studies (14). As the *pmoA* gene phylogeny is largely comparable to the 16S rRNA gene phylogeny (7, 12), *pmoA* primers provide a useful tool for obtaining simultaneous functional and taxonomic inventories of methanotrophs in the environment (10).

Recently, new *pmo*-like (*pxm*) genes in cultured, methanotrophic *Gammaproteobacteria* (21) and new groups of aerobic methane oxidizers with divergent *pmo* genes were discovered. Most noteworthy are the proteobacterial *Crenothrix polyspora* (19) and three acidophilic verrucomicrobial *Methyloirabilis* species (16). The presence of a complete *pmo* gene cluster in the genome of the anaerobic, methanotrophic, nitrite-reducing bacterium “*Candidatus Methyloirabilis oxyfera*” (3) further expanded the array of *pmo* gene diversity. “*Ca. Methyloirabilis oxyfera*,” a member of the uncultured NC10 phylum (17), forms a novel taxonomic group of bacterial methanotrophs (reviewed in reference 24). The presence of *pmo* genes in an anaerobe was explained by the unusual metabolism of “*Ca. Methyloirabilis oxyfera*”: molecular oxygen was produced

from nitric oxide, an intermediate of denitrification, and then used to oxidize methane via the complete aerobic pathway starting with pMMO (3).

It was not possible to amplify *pmoA* genes with the most commonly used forward primer A189 and the reverse primers A682 (7), mb661 (2), and A650 (1), even from “*Ca. Methyloirabilis oxyfera*” enrichment cultures. The alignment of the primers with the *pmoA* sequence of “*Ca. Methyloirabilis oxyfera*” revealed several critical mismatches especially with the reverse primers (Fig. 1), explaining the failure to obtain a PCR product. Based on this alignment, new *pmoA* primers were developed. We changed one nucleotide to a wobble base in the forward primer A189, resulting in a general degenerate primer, A189_b (Fig. 1), matching most methanotrophs in the GenBank database with the exception of the genus *Methyloirabilis*. The reverse primer A682 differed in eight nucleotides from the *pmoA* sequence of “*Ca. Methyloirabilis oxyfera*.” Therefore, the newly developed reverse primer designated cmo682 specifically targets “*Ca. Methyloirabilis oxyfera*”-like bacteria. Also, a second set of primers for a nested-PCR approach specific for nitrite-dependent anaerobic methane-oxidizing bacteria was developed based on this alignment. These primers were named cmo182 and cmo568 (Fig. 1) and were used in a nested approach after the amplification with A189_b and cmo682. The newly designed primers were first tested with DNA extracted from Ooijpolder drainage ditch sediment, which had previously been used to obtain an enrichment culture performing nitrite-dependent anaerobic methane oxidation (4). The same approach was then used to screen samples from several oxygen-limited freshwater habitats for the presence of “*Ca. Methyloirabilis oxyfera*”-like bacteria: an alpine peat bog (China), wastewater treatment plants (WWTP) (Netherlands), and contaminated aquifers (United States and Netherlands) (Table 1). There is circumstantial evidence for the occurrence of anaerobic methane oxidation in the contaminated aquifers (18, 22).

To extract DNA from the samples from Ooijpolder, Banisveld, Rotterdam, and Lichtenvoorde (Table 1), biomass was collected by centrifugation. The pellets were resus-

* Corresponding author. Mailing address: Department of Microbiology, Institute for Water and Wetland Research, Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, Netherlands. Phone: 31243652557. Fax: 31243652830. E-mail: k.ettwig@science.ru.nl.

† Present address: Australian Institute for Bioengineering and Nanotechnology, The University of Queensland, Brisbane, QLD 4072, Australia.

‡ Present address: Max Planck Institute for Marine Microbiology, Celsiusstrasse 1, D-28359 Bremen, Germany.

§ These authors contributed equally to this study.

▽ Published ahead of print on 1 April 2011.

Nucleotide position on <i>M. capsulatus</i> str. Bath	A189b		cmo182		cmo568		cmo682	
	136	153	182	199	549	568	649	666
AE017282 <i>Methylococcus capsulatus</i> str. Bath	ATGGGTGACTGGGACTTCTGGTGG	GGGTACCGGTGACCCGATCGTAC	CATCCAGGGTTACAACTATGTGGTA	TCCGCAATCTTCTCCGCTTCATG				
AB253367 <i>Methylomicrobium japanense</i> str. NI	GGCGGTGACTGGGACTTCTGGACT	GGGTAACTGTAGCGCAATCGTTT	CCTTCAAGGTTACCACTATGTAGAA	TCAGCGTTCTTCTCTGGTTTCGTA				
AB275418 <i>Methylocaldum</i> sp. T 025	ATGGGTGACTGGGACTTCTGGGCG	GGGTACCGGTGACCGGATTTGTGT	CATTCAAGGTTACCACTATGTAGAA	TCGGCGTTCTTCTCGGCTTCATG				
AB501285 <i>Methylovulum miyakonense</i> str. HT12	GGGGACTGGGACTTCTGGGTT	GGGTACCGGTAGTTCGATTTGAT	TATTATGGGTTACCAATACGTCGGA	TCAGCTTTCTTCTCGGTTGGTGA				
DQ119048 <i>Methylosinus sporium</i> str. LC3	GGGGACTGGGACTTCTGGGTC	GGCGACCGGTATCCCGATCCTGG	TCTGATCGGCTCCCACTCGTCGCA	GCCGCTTCTTCTCGGCTTC				
BX649604 <i>Methylocystis</i> str. SC2	GCCGCGGACTGGGACTTCTGGATC	GGCGACCGGTGACCGGATCCTGG	CCTCATCGGCTTCAACTACGTCGCA	GCCGCTTCTTCTCGGCTTCATT				
DQ379514 Uncultured <i>Methylocystis</i> sp. GSC357	GCGGGGACTGGGACTTCTGGGTC	GGCGACCGGTATTTCCGATCCTGG	TCTGATCGGCTTCACTTCTCGTCGCA	GCCGCTTCTTCTCGGCTTCGTC				
EF623812 UB Schoehsee sediment Germany Ssedi43	GGTGACTGGGACTTCTGGGTC	GGGTACCGGTACACCGATTGTAG	TTATATGGGTTACAGTATGTCGTA	TCAGCTTCTTCTCGGCTTC				
EU722433 <i>Methylomonas methanica</i> S1	GGCGATTGGGACTTTTGGGTC	GGCCATTGGTACACCGCTTAATCG	TTTGTTCGGTTTCAATATATCGTA	GCGGCAATTTTCTCGGCACTGTTA				
EF175100 <i>Nitrosospira</i> sp. EnI299	GCAGGCGACTGGGACTTCTGGCTG	GGCGGTAGTGACCCGATCCTGG	CTACACCGGCTTCTGTACGTCGCA	GCCCTATTCTTCTCGGCTTCGTC				
AF042171 <i>Nitrosolobus multififormis</i> str. Nm24C	GCGGGAGACTGGGACTTCTGGCTT	GGCGTGTCTGACCCGATCGTAG	CTACACCGGCTTTTGTATGTAGCA	GCCGCAATCTTCTCGGCTTCGTC				
AL954747 <i>Nitrosomonas europaea</i> str. ATCC 19718	GCAGGTGACTGGGACTTCTGGATG	GGCGGTGTGTAACGCCAATCGTG	TTACATGGGACATCTGTATGTTCGTA	GCAGCAATCTTCTCTCGGCTTCGTA				
DQ295901 <i>Crenothrix polyspora</i> 3	GGACACTGGGACTTCTGGCTC	GGCGATCTGTACACCGATTACCC	CTACATGGGCTTCACTGTATGTTCGTA	TCGGCGTTCTTCTCGGCTTC				
AF047705 <i>Nitrosococcus oceani</i> ATCC19707	ATGGGGGACTGGGACTTCTGGGTT	GGCCACCGGTATTCGATTTGAT	TATTATGGGTTACCAATACGTCGGA	TCAGCTTCTTCTCGGTTTGT				
EF591085 <i>Methylocaldiphilum fumariolicum</i> PmoA1	GCCGGGACTGGGCTTCTGGATC	GGCCAGTGTGTCGCTCATCATGG	CCTGATTCAATATCAGTACATCGGA	TCGGCAATCTTCTCGGCTTCGTC				
EF591086 <i>Methylocaldiphilum fumariolicum</i> PmoA2	GCTGGGATTTGGTCAATTTGGACC	GGCCAGTGTGTCGCTCATCATGG	TCTCATCCAGTATGAATACATCGGA	TCCGCAATCTTCTCGGCTTTGTA				
EF591087 <i>Methylocaldiphilum fumariolicum</i> PmoA3	GAAGGAGACTGGGACTTCTGGATA	TTCCAAATTTAGGAGTATCTCCT	TTACATCGGATTTTCCATGTCGCTT	GCATATTTCTTCAAGTGTGTTGT				
FJ462791 <i>Methylocaldiphilum kamchatkense</i> PmoA4	GCTGGAGACTGGTCTTTTGGATC	GGCCAGTCAATGTTCCAAATTTGCGA	TCTTATTTCAATACAGTATATTCGAA	TCGGCTTCTTCTCGGCTTTATA				
DQ367738 UB Sediment Lake Washington pmoA3	GGGGACTGGGACTTCTGGACC	GGCTGCTGGTGACCGCGTCTCGC	CCTCAGGGTTCCAGTATCATCGCA	TCGGCTTCTTCTCGGCTTC				
EU723734 UB forest soil Hawaii ML22	GGAGACTGGGACTTCTGGACC	GGCTGCTGGTGACCGCGTCTCGC	CTTGCAAGGTTCCAGTATCATCGCA	TCGGCTTCTTCTCGGCTTC				
EU723750 UB forest soil Hawaii CCRd431	GGGGACTGGGACTTCTGGGTC	GGCTGCTGGTGACCGCGTCTCGC	CCTTCAAGGGTTCCAGTATCATCGCA	TCGGCTTCTTCTCGGCTTC				
FP565575 <i>Candidatus Methyloiraxia oxyfera</i>	GTTGGCGACTGGGACTTCTGGGTC	GGGTACCGGTGACCGGATCCTGT	TCTGATGGGGATGAGTATGTCGCT	TCCTCGTCTTCTCGGCGATTATT				
New pmoA primers for "Ca. M. oxyfera"	GGNGACTGGGACTTCTGGGTC	TCACGTTGACGCCGATCC	GATGGGGATGGAGTATGTGC	TCGTTCTTCTCGGCGRTTT				
pmoA primers (Holmes et al. 1995)	GGNGACTGGGACTTCTGGGTC	TCACGTTGACGCCGATCC	GATGGGGATGGAGTATGTGC	GCSTTCTTCTCGGCGRTTT				

FIG. 1. Alignment using MEGA4 of newly developed primers for the alpha subunit of particulate methane monooxygenase (*pmoA*) of "Candidatus Methyloiraxia oxyfera" and nucleotide sequences encoding the *pmoA* genes of different methanotrophic *Gammaproteobacteria* (GenBank accession no. AE017282, AB253367, AB275418, AB501285, DQ295901), methanotrophic *Alphaproteobacteria* (DQ119048, BX649604, DQ379514, EF623812), and *Verrucomicrobia* (EF591085, EF591086, EF591087, FJ462791); different ammonia monooxygenase (*amoA*) sequences of nitrifying *Betaproteobacteria* (EF175100, AF042171, AL954747); *amoA* sequences of nitrifying *Gammaproteobacteria* (AF047705); *pmxA* sequences of methanotrophic *Gammaproteobacteria* (EU722433); and *pmoA* sequences of different uncultured methanotrophs of unknown affiliation (DQ367738, EU723734, EU723750) (20). Identical nucleotides are shaded in gray. cmo568 and cmo682 are shown as the reverse complementary sequence.

pended in 120 mM sodium phosphate buffer, glass beads (~0.3 g, 0.25-mm diameter) were added, and bead beating (Retsch MM301) was performed for 60 s at 30 Hz. DNA was extracted and purified according to reference 13. For the nitrogen-contaminated Cape Cod aquifer (15), 2 liters of groundwater from a depth of 8.9 m (relative to mean sea level) at sampling site F168 was filtered using a Sterivex cartridge filter (0.2- μ m membranes; Millipore). The membrane filter was used for DNA extraction. DNA from Cape Cod and Zoige soil (0.25 g) was extracted using the Power-Soil DNA isolation kit (MO BIO Laboratories, Inc.) according to the manufacturer's protocol.

To amplify the *pmoA* gene fragment, the primers A189_b and cmo682 were used for the first PCR, and cmo182 and cmo568 were used for the second PCR. For both PCRs, thermal cycling was performed with an initial melting step for 4 min at 94°C, followed by 35 cycles of denaturation at 94°C for 1 min, annealing using a temperature gradient from 50 to 60°C (Zoige sample, 53 to 63°C) for 1 min, and elongation at 72°C for 1.5 min. Finally, an elongation step at 72°C for 10 min was performed. After PCR, amplicons of all annealing temperatures were pooled to minimize random polymerase errors and

the effects of primer mismatches. All PCRs were performed with PerfeCTa SYBR Green FastMix (Quanta BioSciences, Inc., Gaithersburg, MD).

PCRs with the primers A189_b and cmo682 on DNA from Ooijpolder sediment yielded faint and multiple PCR bands, whereas the nested PCR using the pooled amplicons of the first PCR with primers cmo182 and cmo568 resulted in a strong and single band on the gel of the correct size (389 bp). The PCR product mix was ligated into the pGEM-T Easy cloning vector and cloned according to the manufacturer's instructions (Promega). Plasmids were isolated from randomly selected clones for each PCR product with the GeneJET miniprep kit (Fermentas, Lithuania). The plasmids were sequenced using the M13 forward primer. Sequences were checked for quality with Chromas Lite 2.01, erroneous sequences were removed, and BLAST search analysis was performed (<http://www.ncbi.nlm.nih.gov/Blast.cgi>). Sequences matching *pmoA* were aligned to a data set of different *pmoA*, *pmxA*, and *amoA* sequences using the MEGA4 software package and the Clustal W algorithm (20). A phylogenetic tree with the *pmoA* sequences of *Methylocaldiphilum* as an outgroup was calculated with the neighbor-joining method, and tree to-

TABLE 1. Overview of sampling locations

Site abbreviation	Sampling location	Geographic coordinate	Environment ^a	Reference
Ooijpolder	Nijmegen, Netherlands	51°50'N, 5°54'E	Ditch	Ettwig et al. (4)
Zoige	Tibetan plateau, China	33°56'N, 102°52'E	Peat bog	Zhang et al. (25)
Cape Cod	Massachusetts, United States	41°41'N, 70°17'W	Aquifer, well F168	Smith et al. (18); Miller and Smith (15)
Banissveld	Oirschot, Netherlands	51°33'N, 5°17'E	Aquifer, 3.2-m depth	van Breukelen and Griffioen (22)
Lichtenvoorde	Lichtenvoorde, Netherlands	51°59'N, 6°34'E	WWTP	WWTP Lichtenvoorde ^b
Rotterdam	Rotterdam, Netherlands	51°55'N, 4°28'E	WWTP	van der Star et al. (23)

^a WWTP, wastewater treatment plant.

^b http://www.waterstromen.nl/onze_activiteiten/afvalwaterbehandeling/afvalwaterzuivering_royal_dutch_tanneries_lichtenvoorde.aspx.




FIG. 2. Phylogenetic tree of *amoA*, *pmoA*, and *pxmA* nucleotide sequences, including the sequences obtained in this study with *Methylacidiphilum* as the outgroup. The tree was calculated using the neighbor-joining method. Bootstrap support values greater than 60% ($n = 1,000$) are indicated at the nodes.

pology robustness was tested by bootstrap analysis of 1,000 replicates (Fig. 2).

Similar to the results with the Ooijpolder samples, the direct PCR on the different environmental samples using the primer combination A189_b and cmo682 yielded multiple PCR products. Therefore, the same nested approach was applied as described above. From all the different environments analyzed (Table 1), *pmoA* sequences closely related to "Ca. Methyloirabilis oxyfera" (at least 85.5% nucleotide identity and 92% protein identity) were retrieved, most of which cluster according to location (Fig. 2). Together they form a distinct group affiliated with the *pmoA* genes of aerobic methanotrophs (Fig. 2). The closest related sequences obtained with PCR using the

conventional *pmoA* primers (8, 9, 11) were not targeted by the new primers (Fig. 1). All newly obtained sequences share the insertion of a proline at position 131 (3), a distinct feature compared to the sequences of known methane or ammonium monooxygenases. The fact that they are not very divergent from each other may reflect functional conservation relative to the oxygenic pathway.

This study provides a rapid and robust method for the amplification of NC10 phylum *pmoA* genes and enables the screening of different environments.

Nucleotide sequence accession numbers. Representative sequences from each location were submitted to GenBank under accession numbers HQ698926 to HQ698937.

We thank Boris van Breukelen and Wilfred Röling for discussions and sampling the Banisveld aquifer, Wiebe Abma (Paques BV) for providing samples from wastewater treatment plants, and Richard L. Smith and Denis R. LeBlanc at the USGS for their support of sampling and field access at the Cape Cod site. We also acknowledge Boran Kartal for critically reading the manuscript.

F.A.L. and T.A.V.A. were supported by the Foundation for Applied Research (STW project no. 07736). B.Z. was supported by the Chinese Academy of Science-Royal Netherlands Academy of Arts and Sciences joint Ph.D. training program. K.F.E. was supported by the ERC (grant no. 2322937) and the Darwin Center for Biogeology. M.S.M.J. was supported by the ERC (grant no. 2322937). M.R.D. is indebted to the Spanish Ministry of Science and Innovation for its support through the PostdExt program. B.S. was supported by the NSF geobiology and low-temperature geochemistry program (EAR-1024900) and the USGS toxic substances hydrology program.

REFERENCES

- Bourne, D. G., I. R. McDonald, and J. C. Murrell. 2001. Comparison of *pmoA* PCR primer sets as tools for investigating methanotroph diversity in three Danish soils. *Appl. Environ. Microbiol.* **67**:3802–3809.
- Costello, A. M., and M. E. Lidstrom. 1999. Molecular characterization of functional and phylogenetic genes from natural populations of methanotrophs in lake sediments. *Appl. Environ. Microbiol.* **65**:5066–5074.
- Ettwig, K. F., et al. 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **464**:543–548.
- Ettwig, K. F., T. van Alen, K. T. van de Pas-Schoonen, M. S. M. Jetten, and M. Strous. 2009. Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum. *Appl. Environ. Microbiol.* **75**:3656–3662.
- Hakemian, A. S., and A. C. Rosenzweig. 2007. The biochemistry of methane oxidation. *Annu. Rev. Biochem.* **76**:223–241.
- Hanson, R. S., and T. E. Hanson. 1996. Methanotrophic bacteria. *Microbiol. Rev.* **60**:439–471.
- Holmes, A. J., A. Costello, M. E. Lidstrom, and J. C. Murrell. 1995. Evidence that particulate methane monooxygenase and ammonia monooxygenase may be evolutionarily related. *FEMS Microbiol. Lett.* **132**:203–208.
- Kalyuzhnaya, M. G., et al. 2006. Fluorescence in situ hybridization-flow cytometry-cell sorting-based method for separation and enrichment of type I and type II methanotroph populations. *Appl. Environ. Microbiol.* **72**:4293–4301.
- King, G. M., and K. Nanba. 2008. Distribution of atmospheric methane oxidation and methanotrophic communities on Hawaiian volcanic deposits and soils. *Microbes Environ.* **23**:326–330.
- Kip, N., et al. 2010. Global prevalence of methane oxidation by symbiotic bacteria in peat-moss ecosystems. *Nat. Geosci.* **3**:617–621.
- Knief, C., S. Kolb, P. L. E. Bodelier, A. Lipski, and P. F. Dunfield. 2006. The active methanotrophic community in hydromorphic soils changes in response to changing methane concentration. *Environ. Microbiol.* **8**:321–333.
- Kolb, S., C. Knief, S. Stubner, and R. Conrad. 2003. Quantitative detection of methanotrophs in soil by novel *pmoA*-targeted real-time PCR assays. *Appl. Environ. Microbiol.* **69**:2423–2429.
- Kowalchuk, G. A., F. J. de Bruijn, I. M. Head, A. D. Akkermans, and J. D. van Elsas. 2004. Molecular microbial ecology manual (MMEM), 2nd ed., vol. 1. Kluwer Academic Publishing, London, United Kingdom.
- McDonald, I. R., L. Bodrossy, Y. Chen, and J. C. Murrell. 2008. Molecular ecology techniques for the study of aerobic methanotrophs. *Appl. Environ. Microbiol.* **74**:1305–1315.
- Miller, D. N., and R. L. Smith. 2009. Microbial characterization of nitrification in a shallow, nitrogen-contaminated aquifer, Cape Cod, Massachusetts and detection of a novel cluster associated with nitrifying *Betaproteobacteria*. *J. Contam. Hydrol.* **103**:182–193.
- Op den Camp, H. J. M., et al. 2009. Environmental, genomic and taxonomic perspectives on methanotrophic *Verrucomicrobia*. *Environ. Microbiol. Rep.* **1**:293–306.
- Raghoebarsing, A. A., et al. 2006. A microbial consortium couples anaerobic methane oxidation to denitrification. *Nature* **440**:918–921.
- Smith, R. L., B. L. Howes, and S. P. Garabedian. 1991. In situ measurement of methane oxidation in groundwater by using natural-gradient tracer tests. *Appl. Environ. Microbiol.* **57**:1997–2004.
- Stoecker, K., et al. 2006. Cohn's *Crenothrix* is a filamentous methane oxidizer with an unusual methane monooxygenase. *Proc. Natl. Acad. Sci. U. S. A.* **103**:2363–2367.
- Tamura, K., J. Dudley, M. Nei, and S. Kumar. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**:1596–1599.
- Tavormina, P. L., V. J. Orphan, M. G. Kalyuzhnaya, M. S. M. Jetten, and M. G. Klotz. 2010. A novel family of functional operons encoding methane/ammonia monooxygenase-related proteins in gamma proteobacterial methanotrophs. *Environ. Microbiol. Rep.* **3**:91–100.
- van Breukelen, B. M., and J. Griffioen. 2004. Biogeochemical processes at the fringe of a landfill leachate pollution plume: potential for dissolved organic carbon, Fe(II), Mn(II), NH₄, and CH₄ oxidation. *J. Contam. Hydrol.* **73**:181–205.
- van der Star, W. R., et al. 2007. Startup of reactors for anoxic ammonium oxidation: experiences from the first full-scale anammox reactor in Rotterdam. *Water Res.* **41**:4149–4163.
- Wu, M. L., et al. 2011. A new intra-aerobic metabolism in the nitrite-dependent anaerobic methane-oxidizing bacterium "*Candidatus* Methylophilus oxyfera." *Biochem. Soc. Trans.* **39**:243–248.
- Zhang, G. S., et al. 2008. Methanogen community in Zoige wetland of Tibetan plateau and phenotypic characterization of a dominant uncultured methanogen cluster ZC-I. *Environ. Microbiol.* **10**:1850–1860.